

Certificate of Mailing

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Mail Stop: AF, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450 on July 21, 2003.

By: [Signature] Printed: Lyza Fimaliar

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: Lal et al.

Title: SERINE DEHYDRATASE HOMOLOG

Serial No.: 09/925,140

Filing Date: August 08, 2001

Examiner: Helms, L.

Group Art Unit: 1642

Mail Stop: AF

Commissioner for Patents

P.O. Box 1450

Alexandria, VA 22313-1450

DECLARATION OF DR. TOD BEDILION
UNDER 37 C.F.R. § 1.132

I, TOD BEDILION, a citizen of the United States, residing at 132 Winding Way, San Carlos, California, declare that:

1. I was employed by Incyte Corporation (hereinafter "Incyte") as a Director of Corporate Development until May 11, 2001. I am currently under contract to be a Consultant to Incyte Corporation.

2. In 1996, I received a Ph.D. degree in Cell, Molecular and Development Biology from UCLA. I had previously received, in 1988, a B.S. degree in biology from UCLA.

Upon my graduation from UCLA, I became, in April 1996, the first employee of Synteni, Inc. (hereinafter "Synteni"). I was a Research Director at Synteni from April 1996 until Synteni was acquired by Incyte in early 1998.

I understand that Synteni was founded in 1994 by T. Dari Shalon while he was a graduate student at Stanford University. I further understand that Synteni was founded for the purpose of commercially exploiting certain "cDNA microarray" technology that was being

worked on at Stanford in the early to mid-1990s. That technology, which I will sometimes refer to herein as the "Stanford-developed cDNA microarray technology", was the subject of Dr. Shalon's doctoral thesis at Stanford. I understand and believe that Dr. P.O. Brown was Dr. Shalon's thesis advisor at Stanford.

During the period beginning before I was employed by Synteni and ending upon its acquisition by Incyte in early 1998, I understand Synteni was the exclusive licensee of the Stanford-developed cDNA microarray technology, subject to any right that the United States government may have with respect to that technology. In early 1998, I understand Incyte acquired rights under the Stanford-developed cDNA microarray technology as part of its acquisition of Synteni.

I understand that at the time of the commencement of my employment at Synteni in April 1996, Synteni's rights with respect to the Stanford-developed cDNA technology included rights under a United States patent application that had been filed June 7, 1995 in the names of Drs. Brown and Shalon and that subsequently issued as United States Patent No. 5,807,522 (the Brown '522 patent). In December 1995, the subject matter of the Brown '522 patent was published based on a PCT patent application that had also been filed in June 1995. The Brown '522 patent (and its corresponding PCT application) describes the use of the Stanford-developed cDNA technology in a number of gene expression monitoring applications, as will be discussed more fully below.

Upon Incyte's acquisition of Synteni, I became employed by Incyte. From early 1998 until late 1999, I was an Associate Research Director at Incyte. In late 1999, I was promoted to the position of Director, Corporate Development.

I have been aware of the Stanford-developed cDNA microarray technology since shortly before I commenced my employment at Synteni. While I was employed by Synteni, virtually all (if not all) of my work efforts (as well as the work efforts of others employed by Synteni) were directed to the further development and commercial exploitation of that cDNA microarray technology. By the end of 1997, those efforts had progressed to the point that I understand Incyte agreed to pay at least about \$80 million to acquire Synteni. Since I have been employed by Incyte, I have continued to work on the further development and commercial

exploitation of the cDNA microarray technology that was first developed at Stanford in the early to mid-1990s.

3. I have reviewed the specification of a United States patent application that I understand was filed on August 8, 2001 in the names of Preeti Lal et al. and was assigned Serial No. 09/925,140 (hereinafter "the Lal '140 application"). Furthermore, I understand that this United States patent application was a divisional application of and claimed priority to United States patent application Serial No. 09/088,435 filed on June 1, 1998 (hereinafter "the Lal '435 application"), having essentially the identical specification, with the exception of corrected typographical errors and reformatting changes. Thus page and line numbers may not match as between the Lal '140 application and the Lal '435 application. In broad overview, the Lal '435 specification pertains to certain nucleotide and amino acid sequences and their use in a number of applications, including gene expression monitoring applications that are useful in connection with (a) developing drugs (e.g., the diagnosis of inherited and acquired genetic disorders, expression profiling, toxicology testing, and drug development with respect to disorders of metabolism and cancer), and (b) monitoring the activity of drugs for purposes relating to evaluating their efficacy and toxicity.

4. I understand that the Lal '140 application contains a claim that recites an isolated polynucleotide selected from the group consisting of a) a polynucleotide comprising the polynucleotide sequence of SEQ ID NO:2 ("hereinafter the "SEQ ID NO:2 polynucleotide"), b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to the polynucleotide sequence of SEQ ID NO:2 over the entire length of SEQ ID NO:2 ("hereinafter "the SEQ ID NO:2 polynucleotide variant"), c) a polynucleotide completely complementary to a polynucleotide of a), d) a polynucleotide completely complementary to a polynucleotide of b), and e) an RNA equivalent of a)-d) (hereinafter "the claimed polynucleotides") and that the Patent Examiner has rejected this claim on the grounds that the specification of the Lal '140 application does not enable the use of the claimed SEQ ID NO:2 polynucleotide variant. I further understand that whether or not a patent specification enables the

use of its claimed subject matter is properly determined from the perspective of a person skilled in the art to which the specification pertains at the time of the patent application was filed.

5. I have been asked (a) to consider with a view to reaching a conclusion (or conclusions) as to whether or not I agree with the Patent Examiner's position that the Lal '140 application and its parent, the Lal '435 application, do not provide an enabling disclosure for the use of the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant, and (b) to state and explain the bases for any conclusions I reach.

6. I have considered the matters set forth in paragraph 5 of this Declaration and have concluded that, contrary to the position I understand the Patent Examiner has taken, the specification of the Lal '435 patent application would have enabled a person skilled in the art at the time of its filing to make and/or use the claimed claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant. More specifically, persons skilled in the art on June 1, 1998 would have understood the Lal '435 application to disclose the use of the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant in a number of gene expression monitoring applications that were well-known at that time to be useful in connection with the development of drugs and the monitoring of the activity of such drugs. I explain the bases for reaching my conclusion in this regard in paragraphs 7-16 below.

7. In reaching the conclusion stated in paragraph 6 of this Declaration, I considered (a) the specification of the Lal '435 application, and (b) a number of published articles and patent documents that evidence gene expression monitoring techniques that were well-known before the June 1, 1998 filing date of the Lal '435 application. The published articles and patent documents I considered are:

(a) Schena, M., Shalon, D., Heller, R., Chai, A., Brown, P.O., and Davis, R.W., Parallel human genome analysis: Microarray-based expression monitoring of 1000 genes, Proc. Natl. Acad. Sci. USA, 93, 10614-10619 (1996) (hereinafter "the Schena 1996 article") (copy annexed at Tab A);

(b) Schena, M., Shalon, D., Davis, R.W., Brown, P.O., Quantitative Monitoring of Gene Expression Patterns with a Complementary DNA Microarray, Science, 270, 467-470 (1995) (hereinafter "the Schena 1995 article") (copy annexed at Tab B);

(c) Shalon and Brown PCT patent application WO 95/35505 titled "Method and Apparatus For Fabricating Microarrays Of Biological Samples," filed on June 16, 1995, and published on December 28, 1995 (hereinafter "the Shalon PCT application") (copy annexed at Tab C);

(d) Brown and Shalon U.S. Patent No. 5,807,522, corresponding to the Shalon PCT application, titled "Methods For Fabricating Microarrays Of Biological Samples," filed on June 7, 1995 and issued on September 15, 1998 (hereinafter "the Brown '522 patent") (copy annexed at Tab D);

(e) DeRisi, J., Penland, L., and Brown, P.O. (Group 1); Bittner, M.L., Meltzer, P.S., Ray, M., Chen, Y., Su, Y.A., and Trent, J.M. (Group 2), Use of a cDNA microarray to analyse gene expression patterns in human cancer, Nat. Genet., 14(4), 457-460 (1996) (hereinafter "the DeRisi article") (copy annexed at Tab E);

(f) Shalon, D., Smith, S.J., and Brown, P.O., A DNA Microarray System for Analyzing Complex DNA Samples Using Two-color Fluorescent Probe Hybridization, Genome Res., 6(7), 639-645 (1996) (hereinafter "the Shalon article") (copy annexed at Tab F);

(g) Heller, R.A., Schena, M., Chai A., Shalon, D., Bedilion, T., Gilmore, J., Woolley, D.E., and Davis R.W., Discovery and analysis of inflammatory disease-related genes using cDNA microarrays, Proc. Natl. Acad. Sci. USA, 94, 2150-2155 (1997) (hereinafter "the Heller article") (copy annexed at Tab G); and

(h) Sambrook, J., Fritsch, E.F., Maniatis, T., Molecular Cloning, A Laboratory Manual, pages 7.37 and 7.38, Cold Spring Harbor Press (1989) (hereinafter "the Sambrook Manual") (copy annexed at Tab H).

8. Many of the published articles and patent documents I considered (i.e., at least items (a)-(f) identified in paragraph 7) relate to work done at Stanford University in the early and mid-1990s with respect to the development of cDNA microarrays for use in gene

expression monitoring applications under which Synteni became exclusively licensed. As I will discuss, a person skilled in the art who read the Lal '435 application on June 1, 1998 would have understood that application to disclose the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant to be useful for a number of gene expression monitoring applications, e.g., as a probe for the expression of that specific polynucleotide in cDNA microarrays of the type first developed at Stanford.

Furthermore, items (a)-(g) establish that gene expression monitoring applications utilizing cDNA microarrays were well-known and established methods routinely used in toxicology testing and drug development at the time of filing the Lal '435 application and for several years prior to June 1, 1998. As such, one of ordinary skill in the art would have recognized that the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant could be used in toxicology testing and drug development, irrespective of the biochemical activities of the encoded polypeptide.

9. Turning more specifically to the Lal '435 specification, the SEQ ID NO:2 polynucleotide is shown at pp. 51-52 as one of two sequences under the heading "Sequence Listing." The Lal '140 specification specifically teaches that the "invention ... provides an isolated and purified polynucleotide comprising the polynucleotide sequence of SEQ ID NO:2" ... (Lal '435 application at p. 3). It further teaches that (a) the identity of the SEQ ID NO:2 polynucleotide was determined from a THP-1 promonocyte cell cDNA library (THP1AZS08) derived from the peripheral blood of a 1-year-old Caucasian male with acute monocytic leukemia (Lal '435 application, p. 14), (b) the SEQ ID NO:2 polynucleotide encodes for the serine dehydratase homolog (SDHH) shown as SEQ ID NO:1 (Lal '435 application at p. 14), and (c) northern analysis of SEQ ID NO:2 shows its expression predominantly in cDNA libraries associated with cancer, inflammation and the immune response, and fetal development and proliferating cell lines (Lal '435 application at p. 14).

The Lal '435 application discusses a number of uses of the in addition to their use in gene expression monitoring applications. However, my discussion in this Declaration concerning the Lal '435 application focuses on the portions of the application that relate to the use of the

claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant in gene expression monitoring applications.

10. The Lal '435 application discloses that the polynucleotide sequences disclosed therein, including the claimed polynucleotides, are useful as probes in microarrays. It further teaches that the microarrays can be used "to monitor the expression level of large numbers of genes simultaneously" for a number of purposes, including "to develop and monitor the activities of therapeutic agents" (Lal '435 application at p. 35, lines 23-26).

In the paragraph immediately following the Lal '435 teachings described in the preceding paragraph of this Declaration, the Lal '435 application teaches that microarrays can be prepared using the previously mentioned cDNA microarray technology developed at Stanford in the early to mid-1990s. In this connection, the Lal '435 application specifically cites to the Schena 1996 article identified in item (a) of paragraph 7 of this Declaration (Lal '435 application at p. 35; *supra*, paragraph 7).

The Schena 1996 article is one of a number of documents that were published prior to the June 1, 1998 priority date of the Lal '435 application that describes the use of the Stanford-developed cDNA technology in a wide range of gene expression monitoring applications, including monitoring and analyzing gene expression patterns in human cancer. In view of the Lal '435 application, the Schena 1996 article, and other related pre-June 1, 1998 publications, persons skilled in the art on June 1, 1998 clearly would have understood the Lal '435 application to disclose the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant to be useful in cDNA microarrays for the development of new drugs and monitoring the activities of drugs for such purposes as evaluating their efficacy and toxicity, as explained more fully in paragraph 15 below.

With specific reference to toxicity evaluations, those of skill in the art who were working on drug development in June 1, 1998 (and for many years prior to June 1, 1998) without any doubt appreciated that the toxicity (or lack of toxicity) of any proposed drug they were working on was one of the most important criteria to be considered and evaluated in connection with the development of the drug. They would have understood at that time that good drugs are not only potent, they are specific. This means that they have strong effects on a specific

biological target and minimal effects on all other biological targets. Ascertaining that a candidate drug affects its intended target, and identification of undesirable secondary effects (i.e., toxic side effects), had been for many years among the main challenges in developing new drugs. The ability to determine which genes are positively affected by a given drug, coupled with the ability to quickly and at the earliest time possible in the drug development process identify drugs that are likely to be toxic because of their undesirable secondary effects, have enormous value in improving the efficiency of the drug discovery process, and are an important and essential part of the development of any new drug. Accordingly, the teachings in the Lal '435 application, in particular regarding use of the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant, in differential gene expression analysis and in the development and the monitoring of the activities of drugs, clearly includes toxicity studies and persons skilled in the art who read the Lal '435 application on June 1, 1998 would have understood that to be so.

11. The Schena 1996 article was not the first publication that described the use of the cDNA microarray technique developed at Stanford to monitor quantitatively gene expression patterns. More than a year earlier (i.e., in October 1995), the Schena 1995 article, titled "Quantitative Monitoring of Gene Expression Patterns with a Complementary DNA Microarray", was published (see Tabs A and B).

12. As previously discussed (*supra*, paragraphs 2 and 7), in the mid-1990s patent applications were filed in the names of Drs. Shalon and Brown that described the Stanford-developed cDNA microarray technology. The two patent documents (i.e., the Shalon PCT application and the Brown '522 patent) annexed to this Declaration at Tabs C and D evidence information that was available to the public regarding the Stanford-developed cDNA microarray technology before the June 1, 1998 filing date of the Lal '435 application.

The Shalon PCT patent application, which was published in December 1995, contains virtually the same (if not exactly the same) specification as the Brown '522 patent. Hence, the Brown '522 patent disclosure was, in effect, available to the public as of the December 1995 publication date of the Shalon PCT application (see Tabs C and D). For the sake of convenience, I cite to and discuss the Brown '522 specification below on the understanding

that the descriptions in that specification were published as of the December 28, 1995 publication date of the Shalon PCT application.

The Brown '522 patent discusses, in detail, the utility of the Stanford-developed cDNA microarrays in gene expression monitoring applications. For example, in the "Summary Of The Invention" section, the Brown '522 patent teaches (see Tab D, col. 4, line 52-col. 5, line 8):

Also forming part of the invention is a method of detecting differential expression of each of a plurality of genes in a first cell type, with respect to expression of the same genes in a second cell type. In practicing the method, there is first produced fluorescent-labeled cDNAs from mRNAs isolated from two cells types, where the cDNAs from the first and second cell types are labeled with first and second different fluorescent reporters.

A mixture of the labeled cDNAs from the two cell types is added to an array of polynucleotides representing a plurality of known genes derived from the two cell types, under conditions that result in hybridization of the cDNAs to complementary-sequence polynucleotides in the array. The array is then examined by fluorescence under fluorescence excitation conditions in which (i) polynucleotides in the array that are hybridized predominantly to cDNAs derived from one of the first or second cell types give a distinct first and second fluorescence emission color, respectively, and (ii) polynucleotides in the array that are hybridized to substantially equal numbers of cDNAs derived from the first and second cell types give a distinct combined fluorescence emission color, respectively. The relative expression of known genes in the two cell types can then be determined by the observed fluorescence emission color of each spot.

The Brown '522 patent further teaches that the "[m]icroarrays of immobilized nucleic acid sequences prepared in accordance with the invention" can be used in "numerous" genetic applications, including "monitoring of gene expression" applications (see Tab D at col. 14, lines 36-42). The Brown '522 patent teaches (a) monitoring gene expression (i) in different tissue types, (ii) in different disease states, and (iii) in response to different drugs, and (b) that arrays disclosed therein may be used in toxicology studies (see Tab D at col. 15, lines 13-18 and 52-58 and col. 18, lines 25-30).

13. Also pertinent to my considerations underlying this Declaration is the DeRisi article, published in December 1996. The DeRisi article describes the use of the Stanford-developed cDNA microarray technology "to analyze gene expression patterns in human cancer" (see Tab E at, e.g., p. 457). The DeRisi article specifically indicates, consistent with what was apparent to persons skilled in the art in December 1996, that increasing the number of genes on the cDNA microarray permits a "more comprehensive survey of gene expression patterns," thereby enhancing the ability of the cDNA microarray to provide "new and useful insights into human biology and a deeper understanding of the gene pathways involved in the pathogenesis of cancer and other diseases" (see Tab E at p. 458).

14. Other pre-June 1, 1998 publications further evidence the utility of the cDNA microarrays first developed at Stanford in a wide range of gene expression monitoring applications (see, e.g., the Shalon and the Heller articles at Tabs F and G). By no later than the March 1997 publication of the Heller article, these publications showed that employees of Synteni (i.e., James Gilmore and myself) had used the cDNA microarrays in specific gene expression monitoring applications (see Tab G).

The Heller article states that the results reported therein "successfully demonstrate the use of the cDNA microarray system as a general approach for dissecting human diseases" (Tab G at p. 2150). Among other things, the Heller article describes the investigation of "1000 human genes that were randomly selected from a peripheral human blood cell library" and "[t]heir differential and quantitative expression analysis in cells of the joint tissue. . . to demonstrate the utility of the microarray method to analyze complex diseases by their pattern of gene expression" (see Tab G at pp. 2150 *et seq.*).

Much of the work reported on in the Heller article was done in 1996. That article, therefore, evidences how persons skilled in the art were readily able, well prior to June 1, 1998, to make and use cDNA microarrays to achieve highly useful results. For example, as reported in the Heller article, a cDNA microarray that was used in some of the highly successful work reported on therein was made from 1,000 genes randomly selected from a human blood cell library.

15. A person skilled in the art on June 1, 1998, who read the Lal '435 application, would understand that application to disclose the claimed polynucleotides, for example, SEQ ID NO:2 or the SEQ ID NO:2 polynucleotide variant, to be highly useful as probes for the expression of that specific polynucleotide in cDNA microarrays of the type first developed at Stanford. For example, the specification of the Lal '435 application would have led a person skilled in the art in June 1, 1998 who was using gene expression monitoring in connection with working on developing new drugs for the treatment of disorders of metabolism and cancer to conclude that a cDNA microarray that contained the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant would be a highly useful tool and to request specifically that any cDNA microarray that was being used for such purposes contain the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant. Persons skilled in the art would appreciate that cDNA microarrays that contained the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant would be a more useful tool than cDNA microarrays that did not contain the polynucleotides in connection with conducting gene expression monitoring studies on proposed (or actual) drugs for treating disorders of metabolism and cancer for such purposes as evaluating their efficacy and toxicity.

I discuss in more detail in items (a)-(f) below a number of reasons why a person skilled in the art, who read the Lal '435 specification in June 1, 1998, would have concluded based on that specification and the state of the art at that time, that the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant would be a highly useful tool for inclusion in cDNA microarrays for evaluating the efficacy and toxicity of proposed drugs for treating disorders of metabolism and cancer, as well as for other evaluations:

(a) The Lal '435 application teaches the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant to be useful as probes in cDNA microarrays of the type first developed at Stanford. It also teaches that such cDNA microarrays are useful in a number of gene expression monitoring applications, including "developing and monitoring the activity of therapeutic agents [i.e., drugs]" (see paragraph 10, *supra*).

(b) By June 1, 1998, the Stanford-developed cDNA microarray technology was a well known and widely accepted tool for use in a wide range of gene expression monitoring applications. This is evidenced, for example, by numerous publications describing

the use of that cDNA technology in gene expression monitoring applications and the fact that, for over a year, the technology had provided the basis for the operations of an up-and-running company (Synteni), with employees, that was created for the purpose of developing and commercially exploiting that technology (see paragraphs 2, 8 and 10-14, *supra*). The fact that Incyte agreed to purchase Synteni in late 1997 for an amount reported to be at least about \$80 million only serves to underscore the substantial practical and commercial significance, in 1997, of the cDNA microarray technology first developed at Stanford (see paragraph 2, *supra*).

(c) The pre-June 1, 1998 publications regarding the cDNA microarray technology first developed at Stanford that I discuss in this Declaration repeatedly confirm that, consistent with the teachings in the Lal '435 application, cDNA microarrays are highly useful tools for conducting gene expression monitoring applications with respect to the development of drugs and the monitoring of their activity. Among other things, those pre-June 1, 1998 publications confirmed that cDNA microarrays (i) were useful for monitoring gene expression responses to different drugs (see paragraph 12, *supra*), (ii) were useful in analyzing gene expression patterns in human cancer, with increasing the number of genes on the cDNA microarray enhancing the ability of the cDNA microarray to provide useful information (see paragraph 13, *supra*), and (iii) were a valuable tool for use as part of a "general approach for dissecting human diseases" and for "analyz[ing] complex diseases by their pattern of gene expression" (see paragraph 14, *supra*).

(d) Based on my own extensive work for a company whose business was the development and commercial exploitation of cDNA microarray technology for more than two years prior to the June 1, 1998 filing date of the Lal '435 application, I have first-hand knowledge concerning the state of the art with respect to making and using cDNA microarrays as of June 1, 1998 (see paragraphs 2 and 14, *supra*). Persons skilled in the art as of that date would have (a) concluded that the Lal '435 application disclosed cDNA microarrays containing the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant to be useful, and (b) readily been able to make and use such microarrays with useful results.

(e) The Lal '435 specification contains a number of teachings that would lead persons skilled in the art on June 1, 1998 to conclude that a cDNA microarray that contained the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant would be a more

useful tool for gene expression monitoring applications relating to drugs for treating disorders of metabolism and cancer than a cDNA microarray that did not contain the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant. Among other things, the Lal '435 specification teaches that the identity of the SEQ ID NO:2 polynucleotide was determined from a THP-1 promonocyte cell cDNA library (THP1AZS08) derived from the peripheral blood of a 1-year-old Caucasian male with acute monocytic leukemia (Lal '435 application, p. 14). Moreover, northern analysis of SEQ ID NO:2 shows its expression predominantly in cDNA libraries associated with cancer, inflammation and the immune response, and fetal development and proliferating cell lines. (Lal '435 application at p. 14). (See paragraph 9, *supra*).

(f) Persons skilled in the art on June 1, 1998 would have appreciated (i) that the gene expression monitoring results obtained using a cDNA microarray containing a probe to a sequence selected from the group consisting of claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant would vary, depending on the particular drug being evaluated, and (ii) that such varying results would occur both with respect to the results obtained from the probe described in (i) and from the cDNA microarray as a whole (including all its other individual probes). These kinds of varying results, depending on the identity of the drug being tested, in no way detracts from my conclusion that persons skilled in the art on June 1, 1998, having read the Lal '435 specification, would specifically request that any cDNA microarray that was being used for conducting gene expression monitoring studies on drugs for treating disorders of metabolism and cancer (*e.g.*, a toxicology study or any efficacy study of the type that typically takes place in connection with the development of a drug) contain any one of the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant as a probe. Persons skilled in the art on June 1, 1998 would have wanted their cDNA microarray to have a probe as described in (i) because a microarray that contained such a probe (as compared to one that did not) would provide more useful results in the kind of gene expression monitoring studies using cDNA microarrays that persons skilled in the art have been doing since well prior to June 1, 1998.

The foregoing is not intended to be an all-inclusive explanation of all my reasons for reaching the conclusions stated in this paragraph 15, and in paragraph 6, *supra*. In my view, however, it provides more than sufficient reasons to justify my conclusions stated in paragraph 6

of this Declaration regarding the Lal '435 application enabling a person skilled in the art at the time of its filing to make and/or use the claimed claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant.

16. Also pertinent to my considerations underlying this Declaration is the fact that the Lal '435 disclosure regarding the uses of the SEQ ID NO:2 polynucleotide for gene expression monitoring applications is not limited to the use of that polynucleotide as a probe in microarrays. For one thing, the Lal '435 disclosure regarding the hybridization technique used in gene expression monitoring applications is broad (Lal '435 application at, e.g., p. 4, lines 19-29).

In addition, the Lal '435 specification repeatedly teaches that the polynucleotides described therein (including the polynucleotide of SEQ ID NO:2) may desirably be used as probes in any of a number of long established "standard" non-microarray techniques, such as Northern analysis, for conducting gene expression monitoring studies. See, e.g.:

(a) Lal '435 application at p. 8, line 30 to p. 9, line1 ("Northern analysis is indicative of the presence of nucleic acids encoding SDHH in a sample, and thereby correlates with expression of the transcript from the polynucleotide encoding SDHH");

(b) Lal '435 application at p. 34, lines 6-9 ("The polynucleotide sequences encoding SDHH may be used in . . . Northern analysis, dot blot, or other membrane-based technologies. . . Such qualitative or quantitative methods are well known in the art");

(c) Lal '435 application at p. 34, lines 20-26 ("In order to provide a basis for the diagnosis of a disorder associated with expression of SDHH, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, . . . with a sequence, or a fragment thereof, encoding SDHH, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used") (emphasis supplied); and

(d) Lal '435 application at p. 41, lines 31-33 ("Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the

hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound (See, e.g., Sambrook, supra, ch. 7.)”

The “Sambrook et al.” reference cited in item (d) immediately above is a reference that was well known to persons skilled in the art in June 1, 1998. A copy of pages from that reference manual, which was published in 1989, is annexed to this Declaration at Tab H. The attached pages from the Sambrook manual provide an overview of northern analysis and other membrane-based technologies for conducting gene expression monitoring studies that were known and used by persons skilled in the art for many years prior to the June 1, 1998 filing date of the Lal ‘435 application.

A person skilled in the art on June 1, 1998, who read the Lal ‘435 specification, would have routinely and readily appreciated that the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant disclosed therein would be useful as a probe to conduct gene expression monitoring analyses using northern analysis or any of the other traditional membrane-based gene expression monitoring techniques that were known and in common use many years prior to the filing of the Lal ‘435 application. For example, a person skilled in the art in June 1, 1998 would have routinely and readily appreciated that the claimed polynucleotides, including the SEQ ID NO:2 polynucleotide variant would be a useful tool in conducting gene expression analyses, using the northern analysis technique, in furtherance of (a) the development of drugs for the treatment of disorders of metabolism and cancer, and (b) analyses of the efficacy and toxicity of such drugs.

Docket No.: PF-0512-1 DIV

17. I declare further that all statements made herein of my own knowledge are true and that all statements made herein on information and belief are believed to be true; and further, that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, and that willful false statements may jeopardize the validity of this application and any patent issuing thereon.



Todd Bedilion

Signed at Redwood City, California
this 21st day of July, 2003